

QM II: Übungszettel 2

(Abgabetermin: 30.04.2014)

Problem 1 - Propagator for the harmonic oscillator (20 pts)

Show that the propagator for the one-dimensional harmonic oscillator is given by

$$K(x, t; x', t_0) = \sqrt{\frac{m\omega}{2\pi i \hbar \sin[\omega(t-t_0)]}} \exp \left[\frac{im\omega}{2\hbar \sin[\omega(t-t_0)]} \{ (x+x')^2 \cos[\omega(t-t_0)] - 2xx' \} \right]$$

Hint: $K(x, t; x', t_0) = \langle x | e^{-\frac{i}{\hbar} \hat{H}(t-t_0)} | x' \rangle$ in general and $\hat{H} = \hat{\mathbf{p}}^2/2m + m\omega \hat{\mathbf{x}}^2/2$ for the harmonic oscillator.

Problem 2 - Electron in a uniform magnetic field (18 pts)

An electron is subject to a uniform time-independent magnetic field of strength B_0 , pointing in the positive z -direction. At $t = 0$, the electron is known to be in an eigenstate of $\hat{\mathbf{S}} \cdot \mathbf{n}$, with eigenvalue $\hbar/2$, where \mathbf{n} is a unit vector, lying in the xz -plane, that makes an angle β with the z -axis.

- Obtain the probability for finding the electron in the $s_x = \hbar/2$ -state as a function of time.
- Find the expectation value of $\hat{\mathbf{S}}_x$ as a function of time.
- Show that your answers make sense in the extreme cases (i) $\beta \rightarrow 0$ and (ii) $\beta \rightarrow \pi/2$.
- Sketch the dynamics on the Bloch sphere. (*You may use matlab, mathematica or a similar program for that.*)

Problem 3 - Heisenberg's equations of motion (12 pts)

The interaction between an electron and a given magnetic field is due to the magnetic moment of the electron,

$$\hat{\boldsymbol{\mu}}_e = -\frac{2e}{m_e c} \hat{\mathbf{S}},$$

and the external magnetic field, \mathbf{B} . The interaction Hamiltonian reads

$$\hat{H} = -\hat{\boldsymbol{\mu}}_e \cdot \mathbf{B}.$$

We consider a polarized electron, with spin polarization (+) in the z -direction, entering a region of constant magnetic field $\mathbf{B} = B_0 \hat{x}$. The electron moves in the y -direction.

- Write the interaction Hamiltonian for this particular problem.
- Solve Heisenberg's equations of motion for the time dependent operators $\hat{\mathbf{S}}_x(t)$, $\hat{\mathbf{S}}_y(t)$ and $\hat{\mathbf{S}}_z(t)$.
- Discuss the results on the Bloch sphere.