

## Kinematically complete measurements of strong field ionization with mid-IR pulses

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2014 J. Phys. B: At. Mol. Opt. Phys. 47 204010

(<http://iopscience.iop.org/0953-4075/47/20/204010>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 141.51.197.187

This content was downloaded on 06/01/2015 at 12:32

Please note that [terms and conditions apply](#).

# Kinematically complete measurements of strong field ionization with mid-IR pulses

Michael G Pullen<sup>1</sup>, Judith Dura<sup>1,6</sup>, Benjamin Wolter<sup>1</sup>,  
Matthias Baudisch<sup>1</sup>, Michaël Hemmer<sup>1</sup>, Nicolas Camus<sup>2</sup>,  
Arne Senftleben<sup>2,3</sup>, Claus Dieter Schroeter<sup>2</sup>, Robert Moshhammer<sup>2</sup>,  
Joachim Ullrich<sup>2,4</sup> and Jens Biegert<sup>1,5</sup>

<sup>1</sup> ICFO-Institut de Ciències Fotòniques, Av Carl Friedrich Gauss 3, E-08860 Castelldefels (Barcelona), Spain

<sup>2</sup> Max-Planck-Institut für Kernphysik, Saupfercheckweg 1, D-69117 Heidelberg, Germany

<sup>3</sup> Institut für Physik und CINsAT, Universität Kassel, Heinrich-Plett-Strasse 40, D-34132 Kassel, Germany

<sup>4</sup> Physikalisch-Technische Bundesanstalt (PTB), Bundesallee 100, D-38116 Braunschweig, Germany

<sup>5</sup> ICREA-Institució Catalana de Recerca i Estudis Avançats, E-08010 Barcelona, Spain

E-mail: [michael.pullen@icfo.eu](mailto:michael.pullen@icfo.eu)

Received 14 March 2014, revised 7 May 2014

Accepted for publication 19 May 2014

Published 8 October 2014

## Abstract

Recent observations of three unique peaks near 1 eV, 100 meV and 1 meV in the electron spectra generated by ionization using intense mid-IR pulses have challenged the current understanding of strong-field (SF) ionization. The results came as a surprise as they could not be reproduced by the standard version of the commonly used SF approximation. We present results showing the simultaneous measurement of all three low energy ranges at high resolution. This capability is possible due to a unique experimental combination of a high repetition rate mid-IR source, which allows probing deep in the quasi-static regime at high data rates, with a reaction microscope, which allows high resolution three dimensional imaging of the electron momentum distribution.

Keywords: optical tunnelling, zero energy structure, low energy structure, OPCPA, mid-IR

(Some figures may appear in colour only in the online journal)


## 1. Strong-field ionization

Strong-field (SF) ionization is the fundamental physical process behind highly nonlinear phenomena such as multi-photon (MP) ionization, above threshold ionization, sequential and non-sequential double ionization, and high harmonic generation. These phenomena are used extensively in modern research topics such as tomographic imaging [1], laser induced electron diffraction [2] and attosecond science [3]. Extraction of physical parameters from the associated experimental observations requires a comparison to theoretical simulations. Fundamental understanding of the SF ionization process is therefore of paramount importance to the continual

progression of these research fields. It is desirable then for theoretical models to qualitatively reproduce the observed features in the electron and ion ionization spectra.

Current understanding of SF ionization is based upon the seminal work of Keldysh from 1964 where ionization induced by an intense electromagnetic wave of low frequency compared to the ionization potential was considered [4]. In this work it was noted that there are two regimes in which ionization can occur: the so-called MP ionization regime, which can be understood by perturbation theory, and the commonly termed tunnelling regime for which perturbation theory fails. In the tunnelling regime the laser electric field can be considered quasi-static (QS) and the interaction approaches the classical limit. Henceforth this region will be called the QS ionization regime. The unitless Keldysh parameter ( $\gamma$ ), which was introduced in [4], provides an eloquent way to distinguish

<sup>6</sup> Present address: Max-Born Institute (MBI), Max-Born Strasse 2A, 12489 Berlin, Germany.



**Figure 1.** The location of the different ionization regimes (a) and the maximum achievable ponderomotive energies  $U_p$  (b) as a function of the peak laser intensity for different laser wavelengths. The blue shaded region in (a) represents the perturbative MP regime while the red shaded region represents the non-perturbative QS limit. The red shaded region in (b) indicates the range over which the laser used in this work operates.

between the two ionization regimes. In atomic units (a.u.), which are used in this paper unless otherwise stated, this parameter is given by,

$$\gamma = \sqrt{\frac{I_p}{2U_p}}, \quad (1)$$

and is a function of the ionization potential ( $I_p$ ) of the target and the cycle averaged ponderomotive energy of the electron in the laser field ( $U_p$ ), which in turn is dependent on the laser peak intensity ( $I$ ), wavelength ( $\lambda$ ) and frequency ( $\omega$ ),

$$U_p = \frac{I\lambda^2}{16\pi^2} = \frac{I}{4\omega^2}. \quad (2)$$


The MP regime is encountered when  $\gamma \gg 1$  while the QS regime is a more appropriate description when  $\gamma \ll 1$ . The two regimes are presented in figure 1(a) for a range of currently achievable laser peak intensities and wavelengths. The presented wavelengths correspond to the ubiquitous Ti:Sapphire technology (0.8  $\mu\text{m}$ ), possible optical parametric amplification (OPA) outputs (1.3 and 2.0  $\mu\text{m}$ ) and the output of the optical parametric chirped pulse amplification (OPCPA) system presented in this work (3.1  $\mu\text{m}$ ) [5]. The transition from MP ionization (blue shading) to QS ionization (red shading) is gradual and a region exists near  $\gamma \sim 1$  where both regimes contribute (white). Most experiments using a wavelength of 0.8  $\mu\text{m}$  operate within this mixed regime.

The benefits of scaling to longer wavelengths are evident in figure 1. Increasing the laser wavelength results in an increased ability to probe deeper in the QS regime. To reach a Keldysh parameter of  $\gamma = 0.4$ , for example, peak intensities of  $\sim 4.3 \times 10^{14}$  W cm<sup>-2</sup> and  $\sim 0.3 \times 10^{14}$  W cm<sup>-2</sup> are required for wavelengths of 0.8  $\mu\text{m}$  and 3.1  $\mu\text{m}$ , respectively. At these intensities, and for a typical pulse duration of  $\sim 50$  fs, Perelomov–Popov–Terent’ev calculations [6] predict that the ground state of Ar would be depleted in the case of 0.8  $\mu\text{m}$  radiation while it would still be unity for a wavelength of 3.1  $\mu\text{m}$ . Thus, the ability to probe deeper into the QS regime by intensity scaling of short wavelength radiation is limited. Wavelength scaling, on the other hand, avoids ionization saturation and so provides a way to probe species with low ionization potentials deep in the QS regime.

Figure 1(b) presents the corresponding increase in achievable electron energies for the same wavelengths used in figure 1(a). With long wavelength sources, ponderomotive energies of hundreds of eV can be obtained. Such electrons can be utilized in laser-induced electron diffraction [2] experiments and to generate soft x-rays via high harmonic generation [7]. The shaded red region indicates the range of ponderomotive energies that the OPCPA system in this paper can generate. To be able to perform both low (1 meV) and high (hundreds of eV) electron energy experiments an electron detection system with a five orders of magnitude dynamic range, such as the reaction microscope (ReMi) used here, is required.

### 1.1. Low energy structure

Up until recently it was believed that the underlying physics of SF ionization were well understood with experimental results predominantly being interpreted in the context of the popular SF approximation (SFA) [4, 8, 9]. With the recent progress in the development of intense mid-IR sources, probing deep into the QS regime has become possible. Unexpected low energy structure (LES) in the electron spectrum were observed using mid-IR wavelengths between 1.7–3.6  $\mu\text{m}$  in combination with a time-of-flight (TOF) spectrometer [10]. The LES was observed near 1 eV and was found to be universal for all target species. It was found that the results presented in [10] were not reproducible by the SFA, which came as a surprise as it was expected to become more accurate as the QS limit was approached. The width of the LES increased with both laser wavelength and intensity, yet was found to be stable for constant  $U_p$ . The observation that it disappeared with circular polarization hinted that its origin might be related to rescattering [11]. Initial calculations [12, 13], which attributed LES to an interplay of multiple forward scattering with a disturbance of the transverse electron momentum by the Coulomb field, indeed supported this proposition. Further investigations attributed the structure to bunching of electrons that miss the ionic core upon recollision (so-called ‘soft recollisions’) [14, 15]. In addition, improved SFA calculations incorporating a second-order rescattering term in the  $S$ -matrix expansion also show agreement with experimental results [16, 17].



**Figure 2.** The complete normalized low energy electron momentum map detected using a ReMi after the ionization of randomly oriented  $N_2$  with the intense  $3.1 \mu\text{m}$  OPCPA based system. A cross-section of the momentum distribution taken at  $p_{||} = 0$  is presented on the right side of the figure where the positions of the LES (green shading), VLES (blue shading) and ZES (red shading) regions in our experiment are also indicated. Typical TOF spectrometers only detect electrons below a  $6^\circ$  acceptance angle (dashed white line). The inset in the bottom right shows the co-ordinate system of the experimental apparatus.

Figure 2 presents a normalized  $N_2$  electron momentum distribution as a function of the initial electron longitudinal ( $p_{||} = p_z$ ) and transverse ( $p_T = \sqrt{p_x^2 + p_y^2}$ ) momenta after ionization by the  $3.1 \mu\text{m}$  OPCPA source described below. The co-ordinate system of the experimental apparatus is shown in the bottom right corner. Gas is injected into the system in the negative  $y$  direction while the  $3.1 \mu\text{m}$  radiation propagates towards positive  $x$ . The laser polarization and the ReMi detection axis are both aligned along  $z$ . The  $y$ -axis of figure 2 has been scaled logarithmically to enhance the visibility of the zero energy structure (ZES) region at low transverse momentum. The white dashed line represents the linear acceptance angle of typical TOF spectrometers and is discussed in more detail in section 2. Its appearance is nonlinear due to the logarithmic scaling of  $p_T$ . A cross-section of the data taken at  $p_{||} = 0$  (dotted vertical line) is presented on the right side of the figure. The transverse momentum range between  $0.1 \text{ a.u.} < p_T < 0.4 \text{ a.u.}$  (shaded green) represents where the majority of the electrons in the LES energy region are detected. The peak near  $p_T = 0.15 \text{ a.u.}$  extends across the range  $-0.55 \text{ a.u.} < p_{||} < 0.55 \text{ a.u.}$  and corresponds to electron energies between  $0.3 \text{ eV} < E < 5 \text{ eV}$ , which agrees well with the energy range in which the LES was observed in [10, 18].

### 1.2. Very low energy structure

Experimental investigations using similar wavelength lasers to [10] revealed so-called very LESs (VLESs) at even lower energies ( $\sim 100 \text{ meV}$ ) than the LES [18]. A ReMi detection system implemented by Wu *et al* provided the ability to measure the initial momentum distributions of the electrons [19]. These results showed a distinctive double hump structure in the VLES longitudinal momentum distribution, which resembled structures observed previously using  $0.8 \mu\text{m}$  radiation [20]. The transverse momentum range in which we predominantly detect electrons corresponding to the VLES region is shown as the shaded blue region roughly between  $0.03 \text{ a.u.} < p_T < 0.1 \text{ a.u.}$  in figure 2.

Theoretical understanding of the VLES is currently not as developed as for the LES. The structure has been attributed

to an influence of the long-range Coulomb potential [19] or a Coulomb interaction at short distances [21], as well as being interpreted as second-order soft recollisions (i.e.: second-order LES) [15, 22]. Recent SFA calculations, which included a second-order rescattering term, failed to reproduce the structure [16]. In addition, for experimental conditions similar to those in which the VLES was first observed, a suppression of the ionization rate due to population trapping in Rydberg states has been reported [23]. Therefore, there seems to be some confusion as to what mechanisms contribute to the electrons detected in this region.

### 1.3. Near zero energy structure


More recently, we have probed even deeper into the QS regime using a high repetition rate  $3.1 \mu\text{m}$  OPCPA based source combined with a ReMi [24]. The ability to achieve Keldysh parameters approaching  $\gamma \sim 0.1$  allowed the observation of a near ZES at  $\sim 1 \text{ meV}$ . The ZES can clearly be observed in figure 2 within the shaded red region between  $0.001 \text{ a.u.} < p_T < 0.03 \text{ a.u.}$ . It is clear that the new structure is outside the detection range of typical TOF spectrometers, which might explain why it avoided observation until recently. The peak was located between 1–10 meV for the studied species and is therefore well separated from the LES or VLES. The results showed that the ZES was as evident as the LES and even more visible than the VLES. Dura *et al* [24] speculated that the origin of the ZES could be related to population trapping in Rydberg states or a rescattering mechanism.

So far no theoretical investigations have been published that reproduce the observed structure, so its exact origin still defies identification. Further high resolution measurements in the deep QS regime, which are currently underway, will provide insight into the problem.

## 2. Experimental apparatus

Previous experimental efforts aimed at investigating structure at low electron energies have predominantly used sources derived from the frequency conversion of Ti:Sapphire radiation





**Figure 3.** An overview of the laser systems that have been used to investigate low energy structures in the deep QS regime. The size of the data points scale logarithmically with the expected ionization yield. The ideal laser would be located in the top-right of the image to ensure QS conditions at high data acquisition rates. Most experiments have been performed with Ti:Sapphire pumped OPA systems and are limited to the bottom left corner. Our 3.1  $\mu\text{m}$  OPCPA system operates at 160 kHz and is therefore perfectly suited for the investigation of low energy structures. The dashed line at 5 kHz represents an estimated future repetition rate limit for intense mid-IR radiation generated using Ti:Sapphire pumped OPA and DFG.


at 0.8  $\mu\text{m}$  via difference frequency generation (DFG) [10] or parametric amplification [10, 18, 19, 23]. A comparison of the laser sources used in the literature is presented in figure 3. Due to the unfavourable wavelength scaling of the ionization rate at constant repetition rate ( $\sim \lambda^{-1.2}$  calculated using an intensity of  $1 \times 10^{14} \text{ W cm}^{-2}$  [25]), the detected yield decreases as one proceeds to probe further into the QS regime. The size of the data points in figure 3 scale logarithmically with the expected yield when both the wavelength scaling and repetition rate are accounted for. The ideal situation would be to be placed in the top right corner of this plot as this is where investigation deep in the QS regime can be achieved with the required high data acquisition rates. The dashed line at 5 kHz represents an estimate of the repetition rate limit for the generation of intense mid-IR radiation using Ti:Sapphire pumped OPA and DFG in the near future.

The majority of the experiments in the literature have operated in the lower left corner of figure 3 using OPA based systems at wavelengths between 1.3–2.3  $\mu\text{m}$ . These sources are typically intensity and repetition rate limited by the 0.8  $\mu\text{m}$  pump. A DFG based investigation with a wavelength of 3.6  $\mu\text{m}$  has been performed [10] but the same intensity and repetition rate limits apply. Dura *et al* [24] recently presented the first results recorded using an OPCPA based system that operated at 3.1  $\mu\text{m}$  and was specifically designed for investigations approaching the QS limit. OPCPA based technology offers many advantages over Ti:Sapphire based mid-IR generation using OPA or DFG. Particularly relevant to this study is the fact that intensities in the  $10^{14} \text{ W cm}^{-2}$  range can be achieved at two orders of magnitude higher repetition rates. As the measured signal can be low when studying nonlinear phenomena, higher repetition rates become vital for keeping measurement times down. Another benefit of our particular OPCPA is that we are not intensity limited as we still have enough pump power available to implement further OPCPA stages if required.

The OPCPA based laser used in this work has been presented previously [5]. Briefly, it provides sub-six-cycle

mid-IR pulses at a repetition rate of 160 kHz. The 3.1  $\mu\text{m}$  radiation is derived from DFG of 1.55 and 1.05  $\mu\text{m}$  pulses in a magnesium oxide doped periodically poled lithium niobate (MgO:PPLN) crystal. It is subsequently chirped and parametrically amplified in four cascaded OPA stages before the pulse is compressed to  $<60$  fs in a grating compressor. After focussing with a 50 mm parabolic mirror peak intensities above  $10^{14} \text{ W cm}^{-2}$  can be achieved, corresponding to Keldysh parameters  $\gamma < 0.3$  for typical atomic and molecular targets. The 3.1  $\mu\text{m}$  radiation is the idler wave in all of the OPCPA stages, which in our case means that it is passively carrier envelope phase (CEP) stable [26]. Combining this capability with few-cycle pulses, which we have already reported [27], opens the possibility of performing CEP resolved measurements of the LES, VLES and ZES in the near future.

In order to take advantage of the deep QS conditions generated by the mid-IR OPCPA we utilize a ReMi detection system. ReMIs possess a number of benefits over typical TOF spectrometers such as: (1) the capability of detecting electrons in coincidence with ions, (2) meV level resolutions at low electron energies and (3) the direct extraction of the full three dimensional electron momentum distribution. The last benefit, in particular, is extremely useful for the investigation of low electron energy structures that have non-negligible transverse momenta [24]. TOF spectrometers generally have a limited maximum acceptance angle of about  $6^\circ$ , as indicated by the dashed white line in figure 2. Only electrons below this line would be detected by a typical TOF spectrometer that has its axis aligned parallel with the laser polarization direction. When only these electrons are included in the analysis, the VLES and ZES regions disappear yet a small peak in the LES energy region is still present. This peak is populated by electrons that originate from just under the acceptance angle limit and therefore most probably correspond to the LES structure presented in [10, 18].



**Figure 4.** The detected electron energy spectra for Ar (a) and  $N_2$  (c) with the different low energy regions indicated by colour. Also shown is the obtained spectra when electrons from all of the low energy regions are included. The corresponding longitudinal momentum distributions are also shown for Ar (b) and  $N_2$  (d).

For more details of the ReMi used in this report see the Methods section of [24] and for a thorough overview of the function and capabilities of ReMIs in general see [28]. Briefly, we inject a cold supersonically expanded gas into vacuum via two skimming stages. The gas is ionized in the interaction region and the resultant charged particles are guided towards opposing position sensitive micro channel plate detectors by homogenous electric and magnetic fields. Momentum distributions of both ions and electrons are then extracted from the position and time of detection. In the case of low electron energy detection the fields are low (an electric field of  $E = 1.5 \text{ V cm}^{-1}$  and a magnetic field of  $B = 4 \text{ G}$ ) to ensure high detection resolutions. A conservative estimate of upper bounds of the longitudinal and transverse momentum resolutions are  $\Delta p_{||} = 0.011 \text{ a.u.}$  and  $\Delta p_T = 0.019 \text{ a.u.}$ , respectively. The fields can also be increased to detect electrons with energies up to hundreds of eV. The capability to resolve electron energies over a range of five orders of magnitude at high data acquisition rates showcases the versatility of this unique apparatus. It is perfectly suited to investigate the kinematics of LESs within the QS regime.


### 3. Results and discussion

We have measured high resolution electron momentum and energy distributions from Ar and isotropically oriented  $N_2$  in the deep QS regime. These targets complement each other with ionization potentials of 15.76 eV and 15.58 eV, respectively. Figure 4(a) presents the electron energy spectrum (linear binning) from Ar detected over four orders of magnitude between 1 meV and  $\sim 4 \text{ eV}$ . The presented electron spectra have an upper energy limit for two reasons. Firstly, the ReMi is configured to provide high resolution imaging at low energy,

which results in an transverse momentum detection limit of  $p_T < 0.4 \text{ a.u.}$ . Secondly, in post-processing we restrict our analysis to  $|p_{||}| < 0.5 \text{ a.u.}$  to avoid false counts introduced by nodal structures from the magnetic field induced electron cyclotron motion.

The black curve in figure 4(a) represents the electron energy spectrum measured for Ar when all electrons within the LES, VLES and ZES momentum regions are analyzed. A broad peak between 0.2–3 eV can be observed that corresponds to the LES presented in [10]. In addition, a narrow peak between 3–10 meV can clearly be discerned. This corresponds to the ZES first presented in [24]. The VLES peak, which has previously been observed near 100 meV [18, 19], cannot directly be observed in the total energy spectrum. However, if the electron selection is limited to the VLES momentum range between 0.02 a.u.  $< p_T < 0.1 \text{ a.u.}$ , as indicated in figure 2, then it becomes more evident (blue curve). The VLES contributions are masked by the LES electrons at higher energies and the ZES electrons at lower energies. Analysis has confirmed that if an acceptance angle limit is enforced during post processing, such as in Wu *et al*, then the VLES peak becomes visible in the electron energy spectrum. The different contributions to the overall spectra for the LES (green curve) and the ZES (red curve) momentum ranges are also presented in figure 4(a).

Further information can be gained by presenting the parallel momentum distributions of the LES, VLES and ZES regions integrated over their respective  $p_T$  limits (as indicated in figure 2). The results of this procedure are presented for Ar in figure 4(b) where each curve has been rescaled between zero and one to highlight the observed features. For reference the result of integrating over the entire  $p_T$  range (all electrons) is also shown. The most striking feature is the ZES peak at  $p_{||} = 0 \text{ a.u.}$  that has a full width at half maximum (FWHM) of



**Figure 5.**  $N_2$  momentum distributions in one perpendicular direction for different parallel momentum ranges are presented in (a). The ranges roughly correspond to the  $p_{||}$  values at which the LES, VLES and ZES regions are observed to start and are graphically indicated in the inset. A Gaussian fit to the ZES curve is presented to highlight the deviation of the cusp-like feature at  $p_y = 0$ . The detected Ar electron energy spectra for linearly and elliptically polarized light is presented in (b). The ZES structure completely disappears for elliptical polarization yet some LES and VLES signal persists.

0.23 a.u. ( $\sim 0.7$  eV). The structure observed in the momentum range attributed to the VLES region drastically contrasts this peak in two ways. Firstly, the peak has split down the middle with a depth that decreases to about 0.8. This feature shows some resemblance to the double peak structure previously reported in [19]. Secondly, the FWHM has increased to 0.48 a.u. (3.1 eV). As the  $p_T$  integration range continues to be increased to values corresponding to the LES region the double hump structure starts to smear out [19]. The FWHM of the LES region has increased to 0.79 a.u. (8.5 eV). When all electrons are included in the distribution, the ZES single peak signal is completely washed out and the VLES double hump structure is significantly suppressed.

The results for the ionization of  $N_2$  are presented in figures 4(c) and (d). The ZES peak in the energy spectrum is at a slightly lower energy of 2 meV while the LES peak is still located near 1 eV. The contributions from the VLES electrons are again obscured in the overall spectrum but the peak near 100 meV becomes more obvious once they are isolated in momentum space. The other noticeable difference between the Ar and  $N_2$  energy spectra is that the LES is comparatively more prominent than the ZES in the case of  $N_2$ . The longitudinal momentum distributions also reveal some other differences between the two targets. Firstly, the widths of the ZES and VLES regions have decreased to FWHMs of 0.12 a.u. (0.2 eV) and 0.30 a.u. (1.2 eV), respectively. Secondly, the VLES double peak depth now reaches to 0.64. Finally, the VLES distribution possesses well defined ‘shoulders’ that are either absent or much less obvious for Ar.

Theoretical investigations into the origins of the LES [12–15, 22] and the VLES [19] have uncovered a wealth of information. However, no published simulations have been able to successfully reproduce the ZES since it was first reported [24]. We are therefore in the unusual position where experimental progress seems to be ahead of current SF ionization understanding. This situation is reminiscent of the initial surprising LES results that Blaga *et al* reported [10]. To help decipher the ZES mystery, we performed additional experiments which may provide some valuable insight into the low energy kinematics that are at play.

Investigation performed by Rudenko *et al* [29] pointed out that cusp-like structures in one of the perpendicular

momentum distributions are unambiguous signatures of long range interactions with the Coulomb field. Figure 5(a) presents the observed normalized  $p_y$  distributions for  $N_2$  over three different  $p_{||}$  integration ranges that roughly correspond to the  $p_{||}$  values at which the LES, VLES and ZES regions are observed to start (green, blue and red curves, respectively). The cusp-like structure can clearly be identified in each region but it is particularly obvious for the ZES region. A Gaussian fit to the ZES data, without including the cusp-like structure, is also presented as the dashed black curve. The fit is excellent for the range  $|p_y| > 0.04$ . These results seem to be universal for all tested targets and they indicate that the Coulomb field is indeed playing a role in the creation of the ZES.

Figure 5(b) presents the energy spectrum of Ar for both linearly and 15% elliptically polarized light. Both curves have been normalized to emphasize their features. The linearly polarized data is similar to that presented in figure 4(a) and shows the familiar ZES and LES peaks. Clearly, the ZES is absent for the elliptically polarized case yet electrons in the VLES and LES regions can still be observed. In fact, a small VLES plateau near 100 meV is visible. If the LES and VLES are interpreted as being due to soft recollisions [15, 22] then the fact that the ZES is absent might point to another mechanism being responsible. It is possible that the ZES is due to population trapping in high lying Rydberg states as it has previously been shown that elliptically polarized light reduces these populations [30]. Further work into the origin is ongoing and will be published at a further date.

## 4. Conclusion

We present kinematically complete three dimensional measurements of the ionization of Ar and  $N_2$  deep in the QS regime. The unique combination of a high repetition rate mid-IR OPCPA source with a ReMi detection system was specifically designed to simultaneously investigate the LES, VLES and the newly discovered ZES regions in both energy and momentum space with unprecedented resolution. The results show that for our experimental conditions the LES, VLES and ZES energy ranges predominantly contain electrons with transverse momenta of 0.1 a.u.  $< p_T < 0.4$  a.u.,



0.03 a.u.  $< p_T < 0.1$  a.u. and  $0.001$  a.u.  $< p_T < 0.03$  a.u., respectively. These three regions also have distinctive features as a function of the longitudinal momentum: the ZES contains a striking peak at  $p_{||} = 0$ , the VLES shows a distinctive double hump structure that possesses a minima at  $p_{||} = 0$ , and the LES region has a more subtle double hump structure with broader wings above  $|p_{||}| > 0.1$ .

We have found that the three low energy regions are present for all atoms and molecules investigated so far (Ar and N<sub>2</sub> in this report and also O<sub>2</sub> in Dura *et al* [24]). The origin of the ZES is still unknown but initial evidence points towards population trapping in high lying states. Further measurements at the high resolutions offered by our system will complement theoretical efforts to track down the responsible mechanism.

## Acknowledgments

We acknowledge support from MINISTERIO DE ECONOMIA Y COMPETITIVIDAD through Plan Nacional (FIS2011-30465-C02-01), the Catalan Agència de Gestió d'Ajuts Universitaris i de Recerca (AGAUR) with SGR 2009-2013, Fundacio Cellex Barcelona, and funding from LASERLAB-EUROPE, grant agreement 228334.

## References

- [1] Itatani J, Levesque J, Zeidler D, Niikura H, Pépin H, Kieffer J C, Corkum P B and Villeneuve D M 2004 Tomographic imaging of molecular orbitals *Nature* **432** 867
- [2] Blaga C I, Xu J, DiChiara A D, Sistrunk E, Zhang K, Agostini P, Miller T A, DiMauro L F and Lin C D 2012 Imaging ultrafast molecular dynamics with laser-induced electron diffraction *Nature* **483** 194
- [3] Krausz F and Ivanov M 2009 Attosecond physics *Rev. Mod. Phys.* **81** 163
- [4] Keldysh L V 1965 Ionization in the field of a strong electromagnetic wave *Sov. Phys.—JETP* **20** 1307
- [5] Thai A, Hemmer M, Bates P K, Chalus O and Biegert J 2011 Sub-250-mrad, passively carrier-envelope-phase-stable mid-infrared OPCPA source at high repetition rate *Opt. Lett.* **36** 3918
- [6] Perelemov A M, Popov V S and Terent'ev M V 1966 Ionization of atoms in an alternating electric field *Sov. Phys.—JETP* **23** 924
- [7] Popmintchev T *et al* 2012 Bright coherent ultrahigh harmonics in the keV x-ray regime from mid-infrared femtosecond lasers *Science* **336** 1287
- [8] Faisal F H M 1973 Multiple absorption of laser photons by atoms *J. Phys. B: At. Mol. Opt. Phys.* **6** L89
- [9] Reiss H R 1980 Effect of an intense electromagnetic field on a weakly bound system *Phys. Rev. A* **22** 1786
- [10] Blaga C I, Catoire F, Colosimo P, Paulus G G, Muller H G, Agostini P and DiMauro L F 2009 Strong-field photoionization revisited *Nature Phys.* **5** 335
- [11] Faisal F H M 2009 Strong-field physics: ionization surprise *Nature Phys.* **5** 319
- [12] Liu C and Hatsagortsyan K Z 2010 Origin of unexpected low energy structure in photoelectron spectra induced by midinfrared strong laser fields *Phys. Rev. Lett.* **105** 113003
- [13] Yan T-M, Popruzhenko S V, Vrakking M J J and Bauer D 2010 Low-energy structures in strong field ionization revealed by quantum orbits *Phys. Rev. Lett.* **105** 253002
- [14] Lemell C, Dimitriou K I, Tong X-M, Nagele S, Kartashov D V, Burgdörfer J and Gräfe S 2012 Low-energy peak structure in strong-field ionization by midinfrared laser pulses: two-dimensional focusing by the atomic potential *Phys. Rev. A* **85** 011403
- [15] Kästner A, Saalmann U and Rost J M 2012 Electron-energy bunching in laser-driven soft recollisions *Phys. Rev. Lett.* **108** 033201
- [16] Guo L *et al* 2013 Scaling of the low-energy structure in above-threshold ionization in the tunneling regime: theory and experiment *Phys. Rev. Lett.* **110** 013001
- [17] Milošević D B 2013 Reexamination of the improved strong-field approximation: low-energy structures in the above-threshold-ionization spectra for short-range potentials *Phys. Rev. A* **88** 023417
- [18] Quan W *et al* 2009 Classical aspects in above-threshold ionization with a midinfrared strong laser field *Phys. Rev. Lett.* **103** 093001
- [19] Wu C Y, Yang Y D, Liu Y Q, Gong Q H, Wu M, Liu X, Hao X L, Li W D, He X T and Chen J 2012 Characteristic spectrum of very low-energy photoelectron from above-threshold ionization in the tunneling regime *Phys. Rev. Lett.* **109** 043001
- [20] Rudenko A, Zrost K, Schröter C D, de Jesus V L B, Feuerstein B, Moshhammer R and Ullrich J 2004 Resonant structures in the low-energy electron continuum for single ionization of atoms in the tunnelling regime *J. Phys. B: At. Mol. Opt. Phys.* **37** L407
- [21] Lin Z-Y, Wu M-Y, Quan W, Lin X-J, Chen J and Cheng Y 2014 The Coulomb effect on a low-energy structure in above-threshold ionization spectra induced by mid-infrared laser pulses *Chin. Phys. B* **23** 023201
- [22] Lemell C, Burgdörfer J, Gräfe S, Dimitriou K I, Arbó D G and Tong X-M 2013 Classical-quantum correspondence in atomic ionization by midinfrared pulses: multiple peak and interference structures *Phys. Rev. A* **87** 013421
- [23] Liu H *et al* 2012 Low yield of near-zero-momentum electrons and partial atomic stabilization in strong-field tunneling ionization *Phys. Rev. Lett.* **109** 093001
- [24] Dura J *et al* 2013 Ionization with low-frequency fields in the tunneling regime *Sci. Rep.* **3** 2675
- [25] Popruzhenko S V, Mur V D, Popov V S and Bauer D 2008 Strong field ionization rate for arbitrary laser frequencies *Phys. Rev. Lett.* **101** 193003
- [26] Baltuška A, Fujii T and Kobayashi T 2002 Controlling the carrier-envelope phase of ultrashort light pulses with optical parametric amplifiers *Phys. Rev. Lett.* **88** 133901
- [27] Hemmer M, Baudisch M, Thai A, Couairon A and Biegert J 2013 Self-compression to sub-3-cycle duration of mid-infrared optical pulses in dielectrics *Opt. Express* **21** 28095
- [28] Ullrich J, Moshhammer R, Dorn A, Dörner R, Schmidt L P H and Schmidt-Böcking H 2003 Recoil-ion and electron momentum spectroscopy: reaction-microscopes *Rep. Prog. Phys.* **66** 1463
- [29] Rudenko A, Zrost K, Ergler T, Voitkiv A B, Najjari B, de Jesus V L B, Feuerstein B, Schröter C D, Moshhammer R and Ullrich J 2005 Coulomb singularity in the transverse momentum distribution for strong-field single ionization *J. Phys. B: At. Mol. Opt. Phys.* **38** L191
- [30] Nubbemeyer T, Gorling K, Saenz A, Eichmann U and Sandner W 2008 Strong-field tunneling without ionization *Phys. Rev. Lett.* **101** 233001